



Fig 1A




FIG. 1A

Title: IMPROVED CONDITIONALLY REPLICATING
VECTORS FOR INHIBITING VIRAL INFECTION


Inventor: Laurent HUMEAU et al

Application No.: 09/819,401 - Docket No. 397272005/00

Sheet 2 of 49




Fig 1B




09819401-001001



Fig 1C




09819401.091001



09819401.093001



Fig 1E



09819401-091001



Fig 1F






Fig 16






Fig 1H






Fig 1 I



09819401-0001




Fig 1J



09819401-091001



Fig 1K



09819401-091001



A +105 GTGTGCCCGTCTG +117
B AC . . .

A +118 TTGTGTGACTCTG +130
B

A +131 GTAAGTAGAGATC +143
B . C . G A .


FIG. 2

0919101-091001

FIGURE 13

3A

Ratio Optimization for pN1(cPTC)ASenvGFP Vector






TOOT50" T046T860

24

Ratio Optimization for pN1(cPT)GFP Vectors

Title: IMPROVED CONDITIONALLY REPLICATING
VECTORS FOR INHIBITING VIRAL INFECTIONS
First Inventor: Laurent HUMEAU et al
Application No.: 09/819,401 - Docket No. 397272000700
Sheet 14 of 49






FOOTNOTES

3C

Ratio Optimization for pN1(cPT2)ASenvGFP Vector






FOOTNOTES

3b

Best Vector to Packaging Ratio for pN1cGFP Vector

Title: IMPROVED CONDITIONALLY REPLICATING
VECTORS FOR INHIBITING VIRAL INFECTIONS
First Inventor: Laurent HUÉAU et al
Application No.: 09/819,401 - Docket No. 397272000700
Sheet 16 of 49





FOOTED" T046T860

3E

Optimization of vector to packaging ratio for pN2cGFP

Title: IMPROVED CONDITIONALLY REPLICATING
VECTORS FOR INHIBITING VIRAL INFECTIONS
First Inventor: Laurent HUMEAU et al
Application No.: 09/819,401 - Docket No. 397272000700
Sheet 17 of 49






Fig 4A



0931601046T860

Fig 4B






FOOTNOTES


5A


Ratio Optimization for Packaging of pS1cGFP vectors.

Title: IMPROVED CONDITIONALLY REPLICATING
VECTORS FOR INHIBITING VIRAL INFECTIONS
First Inventor: Laurent HUMEAU et al
Application No.: 09/819,401 - Docket No. 397272000700
Sheet 20 of 49




Optimization of vector to packaging ratio for pS2cGFP





Packaging Construct



New features:

- First 42 nt of gag are degenerated.
- Tat and rev represented as cDNA.
- First 208 nt of rev and last 183 nt of tat are degenerated.
- RRE from HIV-2 is used instead of HIV-1 RRE.


These features eliminate almost any homology with the vector plasmid, make system safer.

- Anti-U5 ribozyme is expressed within gag/pol/RRE cassette, further improving safety.
- Gag/pol/rev/tat/RRE cassette and VSV-G expressed from the same plasmid. This feature may enhance packaging efficiency and titers of the vectors.

09819401-091001



**Fig. 6B Packaging Plasmid
for Second Generation
Vectors**



09819401-091001






Fig. 6C Packaging Plasmid
for First Generation Vectors




09819401.031001

Fig 6 D



09819401-051001

Fig 6E



09819401.03.001

Fig 6F





Fig 66





09819401-01001

FIG. 1

Fig

Influence of Ribozyme(s) in the Packaging on pN1(cPT)GFP Vector Titers in HeLa-tat Cells






TOOT60" T046T860

Fig 8

Challenge #26, MOI 0.1, 100% transduced






Figure 9A

FOOT 60" T046T860




Figure 9B

Potent Inhibition of Wild-type HIV Replication
 by Smartvector Containing T Cells





FIG. 10A


Fig 10A



T00T60" T046T860

10.5

Expansion of SupT1 cells after BG & BCNU














Fig 12



Marker

1 pN1 CGFP 1C exp 30

3 pN1 CGFP 2C exp 30

1-4 pVP1.2

9-12 pVP1.2 Rz

13-16 pVP1.2 Rz2

pNL4-3 with DNase I

pNL4-3 without DNase I


Amp. Neg. Control

Extraction Neg. Control

Marker

100T60" T046T860

Fig 13A



Histogram Statistics

File: SupT1 exp.52 transd..009
Tube: pN1(cPT)ASenvGFP 452 a

Sample ID: SupT1 e
Acquisition Date: 25-

Marker	Left, Right	Events	% Gated	% Total	Mean
All	1, 9910	6356	100.00	63.56	570.39
M1	1, 20	95	1.49	0.95	13.86
M2	20, 9910	6262	98.52	62.62	578.74

0981401-0981401






Fig 14A

Vsv-G, RD114 AND RD114-VSV-G CHIMERIA ENVELOPE PROTEINS




Fig 14B


Titers of RD114-pseudotyped HIV-1 vectors in HT1080

Envelopes	IU/ml
VSV G	3.5x10e6
Rabies virus G	1.6x10e6
RD114WT env	1.5x10e5
RD114E env	3.8x10e4

09819401.03001




Fig 15A



09819401.091001




Fig 15B



09819401-091001



Fig 15C



09819401.091001



Fig 15D





FIG. 15D

Fig 15 E



0561051051051

Fig 15f

Rev dependent VSV-G constructs




Figure 2



Yield of pN1(cPT)GFP Vectors per Cell Factory before and after Concentration in HeLa-tat Cells.




Fig 16


AFTER
CONCENTRATION

BEFORE
CONCENTRATION



0914010360

July 17



TETRACYCLINE	+	-	+	-	+	-	+	-	+	-	+	-
	X PCMV-REV		3 PCMV-REV		5 PCMV-REV		4 PCMV-REV		9 PCMV-REV		+ PCMV-REV	
293G												


- + = PCMV-REV
- = PCI
- G = β -globin SD
- IM-HIV-1 major SD
- H-Haymanarsky's SD
Analog
- IE-HIV-1 env SD
- 2E-HIV-2 env SD

REMOVE TETRACYCLINE
TO INDUCE EXPRESSION OF VSV-G
THAT IS ~~DEPENDENT~~ DEPENDENT.



F.2 18

Influence of the Buffer on Vector Recovery after Storage for 3-5 Weeks at Different Temperatures




- D-PBS
- D-PBS+10% trehalose 1:1
- D-PBS+10% mannitol 1:1
- D-PBS+10% glucose 1:1
- D-PBS+10% trehalose 1:1
- D-PBS+10% glucose 1:1
- D-PBS+10% mannitol 1:1
- D-PBS+10% glucose 1:1
- D-PBS+10% mannitol 1:1
- D-PBS+10% glucose 1:1





Figure 19



09/819,401 - Docket No. 397272000700